

SCALING AND SCORING OF THE

Pediatric Quality of Life Inventory™

PedsQL™

Mapi Research Trust
27 rue de la Villette
69003 Lyon
France
Phone: +33 (0) 4 72 13 65 75
Fax: +33 (0) 4 72 13 66 82
Email: PROinformation@mapi-trust.org

James W. Varni, Ph.D.
Professor
Department of Pediatrics, College of Medicine
Professor, Department of Landscape
Architecture and Urban Planning, College of
Architecture
3137 TAMU
College Station, Texas 77843-3137
USA
Email: jvarni@arch.tamu.edu

PedsQL™ 4.0 Generic Core Scales	5
PARENT Report for Toddlers (ages 2-4)	5
PedsQL™ 4.0 Generic Core Scales	7
CHILD and PARENT Reports for Young Children (ages 5-7), Children (ages 8-12), Teens (ages 13-18)	7
PedsQL™ 4.0 Generic Core Scales	9
YOUNG ADULT Report (ages 18-25) and ADULT Report (ages over 26).....	9
PedsQL™ Infant Scales	11
PARENT Report for Infants (ages 1-24 months).....	11
PedsQL™ 4.0 SF15 Short Form Generic Core Scales	14
PARENT Report for Toddlers (ages 2-4) CHILD and PARENT Reports for Young Children (ages 5-7), Children (ages 8-12), Teens (ages 13-18).....	14
PedsQL™ General Well-Being Scale	17
CHILD, ADOLESCENT and YOUNG ADULT Report (ages 8-25 years of age)	17
PedsQL™ 3.0 Arthritis Module	19
PARENT Report for Toddlers (ages 2-4)	19
PedsQL™ 3.0 Arthritis Module	21
CHILD and PARENT Reports for Young Children (ages 5-7)	21
PedsQL™ 3.0 Arthritis Module	23
CHILD and PARENT Reports for Children (ages 8-12), Teens (ages 13-18)	23
PedsQL™ 3.0 Asthma Module	25
PARENT Report for Toddlers (ages 2-4)	25
PedsQL™ 3.0 Asthma Module	27
CHILD and PARENT Reports for Young Children (ages 5-7), Children (ages 8-12), Teens (ages 13-18).....	27
PedsQL™ 3.0 SF22 Short Form Asthma Module	29
PARENT Report for Toddlers (ages 2-4)	29
PedsQL™ 3.0 SF22 Short Form Asthma Module	31
CHILD and PARENT Reports for Young Children (ages 5-7), Children (ages 8-12), Teens (ages 13-18).....	31
PedsQL™ Brain Tumor Module	33
PARENT Report for Toddlers (ages 2-4)	33
PedsQL™ Brain Tumor Module	35
CHILD and PARENT Reports for Young Children (ages 5-7)	35
PedsQL™ Brain Tumor Module	37
CHILD and PARENT Reports for Children (ages 8-12), Teens (ages 13-18)	37
PedsQL™ 3.0 Cancer Module	39
PARENT Report for Toddlers (ages 2-4)	39
PedsQL™ 3.0 Cancer Module	41
CHILD and PARENT Reports for Young Children (ages 5-7)	41
PedsQL™ 3.0 Cancer Module	43
CHILD and PARENT Reports for Children (ages 8-12), and Teens (ages 13-18), and Young Adults Report (ages 18-25)	43
PedsQL™ 3.0 Cardiac Module	45
PARENT Report for Toddlers (ages 2-4)	45
PedsQL™ 3.0 Cardiac Module	47
CHILD and PARENT Reports for Young Children (ages 5-7)	47
PedsQL™ 3.0 Cardiac Module	49

CHILD and PARENT Reports for Children (ages 8-12) and Teens (ages 13-18)	49
PedsQL™ 3.0 Cerebral Palsy Module	51
PARENT Report for Toddlers (ages 2-4)	51
PedsQL™ 3.0 Cerebral Palsy Module	53
CHILD and PARENT Reports for Young Children (ages 5-7), Children (ages 8-12), Teens (ages 13-18)	53
PedsQL™ Cognitive Functioning Scale.....	55
PedsQL™ 3.0 Diabetes Module.....	57
PedsQL™ 3.2 Diabetes Module.....	59
PedsQL™ 3.0 Duchenne Muscular Dystrophy Module.....	61
PARENT Report for Young Children (ages 5-7).....	61
PedsQL™ 3.0 Duchenne Muscular Dystrophy Module.....	63
CHILD and PARENT Reports for Children (ages 8-12) and Teens (ages 13-18)	63
PedsQL™ 3.0 Eosinophilic Esophagitis Module.....	65
PARENT Report for Toddlers (ages 2-4)	65
PedsQL™ 3.0 Eosinophilic Esophagitis Module.....	68
CHILD and PARENT Reports for Young Children (ages 5-7)	68
PedsQL™ 3.0 Eosinophilic Esophagitis Module.....	71
CHILD and PARENT Reports for Children (ages 8-12) and Teens (ages 13-18)	71
PedsQL™ 3.0 End Stage Renal Disease Module	74
PARENT Report for Toddlers (ages 2-4)	74
PedsQL™ 3.0 End Stage Renal Disease Module	76
CHILD and PARENT Reports for Young Children (ages 5-7), Children (ages 8-12), Teens (ages 13-18).....	76
PedsQL™ 2.0 Family Impact Module	78
PARENT Report	78
PedsQL™ Gastrointestinal Symptom Scale (GSS)	81
PARENT Proxy Report.....	81
PedsQL™ Healthcare Satisfaction Generic Module.....	83
PedsQL™ Healthcare Satisfaction Hematology / Oncology Module.....	85
PedsQL™ Multidimensional Fatigue Scale.....	87
PedsQL™ 1.0 Neurofibromatosis Type 1 Module.....	89
PedsQL™ 3.0 Neuromuscular Module.....	92
PedsQL™ Oral Health Scale	94
PedsQL™ Pediatric Pain Coping Inventory™ (PPCI).....	96
PARENT Report for Children and Teens (ages 5-18)	96
CHILD Report for Children (ages 5-12), Teens (ages 13-18)	96
PedsQL™ Pediatric Pain Questionnaire™ (PPQ)	98
PedsQL™ Present Functioning Visual Analogue Scale (PedsQL™ VAS)	100
CHILD and PARENT reports for CHILD (ages 5-18)	100
PedsQL™ 3.0 Rheumatology Module.....	102
PARENT Report for Toddlers (ages 2-4)	102
PedsQL™ 3.0 Rheumatology Module.....	104
CHILD and PARENT reports for Young Children (ages 5-7)	104
PedsQL™ 3.0 Rheumatology Module.....	106
CHILD and PARENT reports for Children (ages 8-12) and Teens (ages 13-18).....	106
PedsQL™ 3.0 Sickle Cell Disease Module.....	108
PARENT Report for Toddlers (ages 2-4)	108
PedsQL™ 3.0 Sickle Cell Disease Module.....	110

CHILD and PARENT Reports for Young Children (ages 5-7).....	110
PedsQL™ 3.0 Sickle Cell Disease Module.....	112
CHILD and PARENT Reports for Children (ages 8-12) and Teens (ages 13-18)	112
PedsQL™ 3.0 Transplant Module	114

PedsQL[™] 4.0 Generic Core Scales

PARENT Report for Toddlers (ages 2-4)

The **Parent Report for Toddlers** (ages 2-4) of the **PedsQL™ 4.0 Generic Core Scales** is composed of 21 items comprising 4 dimensions.

DESCRIPTION OF THE QUESTIONNAIRE:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
Physical Functioning	8	1-8	1-8	Higher scores indicate better HRQOL.
Emotional Functioning	5	1-5	1-5	
Social Functioning	5	1-5	1-5	
School Functioning	3	1-3	1-3	

SCORING OF DIMENSIONS:

Item Scaling	5-point Likert scale from 0 (Never) to 4 (Almost always)
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed on a scale from 0 to 100.
Scoring Procedure	<p><u>Step 1: Transform Score</u></p> <p>Items are reversed scored and linearly transformed to a 0-100 scale as follows: 0=100, 1=75, 2=50, 3=25, 4=0.</p> <p><u>Step 2: Calculate Scores</u></p> <p><u>Score by Dimensions:</u></p> <ul style="list-style-type: none"> If more than 50% of the items in the scale are missing, the scale scores should not be computed. Mean score = Sum of the items over the number of items answered. <p><u>Psychosocial Health Summary Score</u> = Sum of the items over the number of items answered in the Emotional, Social, and School Functioning Scales.</p> <p><u>Physical Health Summary Score</u> = Physical Functioning Scale Score</p> <p><u>Total Score:</u> Sum of all the items over the number of items answered on all the Scales.</p>
Interpretation and Analysis of Missing Data	<p>If more than 50% of the items in the scale are missing, the Scale Scores should not be computed.</p> <p>If 50% or more items are completed: Impute the mean of the completed items in a scale.</p>

PedsQL[™] 4.0 Generic Core Scales

**CHILD and PARENT Reports for Young Children (ages 5-7),
Children (ages 8-12), Teens (ages 13-18)**

The **Child and Parent Reports** of the **PedsQL™ 4.0 Generic Core Scales** for:

- Young Children (ages 5-7),
- Children (ages 8-12),
- And Teens (ages 13-18),

are composed of 23 items comprising 4 dimensions.

DESCRIPTION OF THE QUESTIONNAIRE:

Dimensions	Number of Items	Cluster of Items	Reversed scoring	Direction of Dimensions
Physical Functioning	8	1-8	1-8	Higher scores indicate better HRQOL.
Emotional Functioning	5	1-5	1-5	
Social Functioning	5	1-5	1-5	
School Functioning	5	1-5	1-5	

SCORING OF DIMENSIONS:

Item Scaling	5-point Likert scale from 0 (Never) to 4 (Almost always) 3-point scale: 0 (Not at all), 2 (Sometimes) and 4 (A lot) for the Young Child (ages 5-7) child report
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed on a scale from 0 to 100.
Scoring Procedure	<p><u>Step 1: Transform Score</u></p> <p>Items are reversed scored and linearly transformed to a 0-100 scale as follows: 0=100, 1=75, 2=50, 3=25, 4=0.</p> <p><u>Step 2: Calculate Scores</u></p> <p><u>Score by Dimensions:</u></p> <ul style="list-style-type: none"> • If more than 50% of the items in the scale are missing, the scale scores should not be computed, • Mean score = Sum of the items over the number of items answered. <p><u>Psychosocial Health Summary Score</u> = Sum of the items over the number of items answered in the Emotional, Social, and School Functioning Scales.</p> <p><u>Physical Health Summary Score</u> = Physical Functioning Scale Score</p> <p><u>Total Score:</u> Sum of all the items over the number of items answered on all the Scales.</p>
Interpretation and Analysis of Missing Data	If more than 50% of the items in the scale are missing, the Scale Scores should not be computed. If 50% or more items are completed: Impute the mean of the completed items in a scale.

PedsQL[™] 4.0 Generic Core Scales

**YOUNG ADULT Report (ages 18-25) and ADULT Report (ages
over 26)**

The **Young Adult and Adult Reports** of the **PedsQL™ 4.0 Generic Core Scales** for:

- Young Adults (ages 18-25),
- Adults (ages over 26),

are composed of 23 items comprising 4 dimensions.

DESCRIPTION OF THE QUESTIONNAIRE:

Dimensions	Number of Items	Cluster of Items	Reversed scoring	Direction of Dimensions
Physical Functioning	8	1-8	1-8	Higher scores indicate better HRQOL.
Emotional Functioning	5	1-5	1-5	
Social Functioning	5	1-5	1-5	
School Functioning	5	1-5	1-5	

SCORING OF DIMENSIONS:

Item Scaling	5-point Likert scale from 0 (Never) to 4 (Almost always)
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed on a scale from 0 to 100.
Scoring Procedure	<p><u>Step 1: Transform Score</u></p> <p>Items are reversed scored and linearly transformed to a 0-100 scale as follows: 0=100, 1=75, 2=50, 3=25, 4=0.</p> <p><u>Step 2: Calculate Scores</u></p> <p><u>Score by Dimensions:</u></p> <ul style="list-style-type: none"> • If more than 50% of the items in the scale are missing, the scale scores should not be computed, • Mean score = Sum of the items over the number of items answered. <p><u>Psychosocial Health Summary Score</u> = Sum of the items over the number of items answered in the Emotional, Social, and School Functioning Scales.</p> <p><u>Physical Health Summary Score</u> = Physical Functioning Scale Score</p> <p><u>Total Score:</u> Sum of all the items over the number of items answered on all the Scales.</p>
Interpretation and Analysis of Missing Data	<p>If more than 50% of the items in the scale are missing, the Scale Scores should not be computed.</p> <p>If 50% or more items are completed: Impute the mean of the completed items in a scale.</p>

PedsQL[™] Infant Scales

PARENT Report for Infants (ages 1-24 months)

The Parent Report for Infants (ages 1-24 months) of the **PedsQL™ Infant Scales** is composed of 36 items for infants ages 1-12 months and 45 items for infants ages 13-24 months comprising 5 dimensions.

DESCRIPTION OF THE QUESTIONNAIRE (INFANTS 1-12 MONTHS):

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
Physical Functioning	6	1-6	1-6	Higher scores indicate better HRQOL.
Physical Symptoms	10	1-10	1-10	
Emotional Functioning	12	1-12	1-12	
Social Functioning	4	1-4	1-4	
Cognitive Functioning	4	1-4	1-4	

DESCRIPTION OF THE QUESTIONNAIRE (INFANTS 13-24 MONTHS):

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
Physical Functioning	9	1-9	1-9	Higher scores indicate better HRQOL.
Physical Symptoms	10	1-10	1-10	
Emotional Functioning	12	1-12	1-12	
Social Functioning	5	1-5	1-5	
Cognitive Functioning	9	1-9	1-9	

SCORING OF DIMENSIONS:

Item Scaling	5-point Likert scale from 0 (Never) to 4 (Almost always)
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed on a scale from 0 to 100.

<p>Scoring Procedure</p>	<p><u>Step 1: Transform Score</u></p> <p>Items are reversed scored and linearly transformed to a 0-100 scale as follows: 0=100, 1=75, 2=50, 3=25, 4=0.</p> <p><u>Step 2: Calculate Scores</u></p> <p><u>Score by Dimensions:</u></p> <ul style="list-style-type: none"> • If more than 50% of the items in the scale are missing, the scale scores should not be computed. • Mean score = Sum of the items over the number of items answered. <p><u>Psychosocial Health Summary Score</u> = Sum of the items over the number of items answered in the Emotional, Social, and Cognitive Functioning Scales.</p> <p><u>Physical Health Summary Score</u> = Sum of the items over the number of items answered in the Physical Functioning and Physical Symptoms Scales.</p> <p><u>Total Score:</u> Sum of all the items over the number of items answered on all the Scales.</p>
<p>Interpretation and Analysis of Missing Data</p>	<p>If more than 50% of the items in the scale are missing, the Scale Scores should not be computed.</p> <p>If 50% or more items are completed: Impute the mean of the completed items in a scale.</p>

PedsQL™ 4.0 SF15

Short Form Generic Core Scales

**PARENT Report for Toddlers (ages 2-4)
CHILD and PARENT Reports for Young Children (ages 5-7),
Children (ages 8-12), Teens (ages 13-18)**

The **Parent Report for Toddlers** (ages 2-4) and the **Child & Parent Reports for Young Children** (ages 5-7), **Children** (ages 8-12) and **Teens** (ages 13-18) of the **PedsQL™ 4.0 SF15 Generic Core Scales** are composed of 15 items comprising 4 dimensions.

DESCRIPTION OF THE SF15 QUESTIONNAIRE:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
Physical Functioning	5	1-5	1-5	Higher scores indicate better HRQOL.
Emotional Functioning	4	1-4	1-4	
Social Functioning	3	1-3	1-3	
School Functioning	3	1-3	1-3	

SCORING OF DIMENSIONS:

Item Scaling	5-point Likert scale from 0 (Never) to 4 (Almost always) 3-point scales: 0 (Not at all), 2 (Sometimes) and 4 (A lot) for the Young Child self-report
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed on a scale from 0 to 100.
Scoring Procedure	<p><u>Step 1: Transform Score</u></p> <p>Items are reversed scored and linearly transformed to a 0-100 scale as follows: 0=100, 1=75, 2=50, 3=25, 4=0.</p> <p><u>Step 2: Calculate Scores</u></p> <p><u>Score by Dimensions:</u></p> <ul style="list-style-type: none"> If more than 50% of the items in the scale are missing, the scale scores should not be computed, Mean score = Sum of the items over the number of items answered. <p><u>Psychosocial Health Summary Score</u> = Sum of the items over the number of items answered in the Emotional, Social, and School Functioning Scales.</p> <p><u>Physical Health Summary Score</u> = Physical Functioning Scale Score</p> <p><u>Total Score:</u> Sum of all the items over the number of items answered on all the Scales.</p>

Interpretation and Analysis of Missing Data	If more than 50% of the items in the scale are missing, the Scale Scores should not be computed. If 50% or more items are completed: Impute the mean of the completed items in a scale.
--	--

PedsQL[™] General Well-Being Scale

**CHILD, ADOLESCENT and YOUNG ADULT Report (ages 8-25
years of age)**

The **PedsQL™ General Well-Being Scale** is composed of 6 items for ages 8-25 years of age, and 1 overall General Health item.

DESCRIPTION OF THE QUESTIONNAIRE (ages 8-25 years):

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
General Well-Being	6	1-6	No	Higher scores indicate better HRQOL.

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
General Health	1	1	No	Higher scores indicate better HRQOL.

SCORING OF DIMENSIONS:

Item Scaling	5-point Likert scale from 0 (Never) to 4 (Almost always)
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed on a scale from 0 to 100.
Scoring Procedure	<p><u>Step 1: Transform Score</u></p> <p>Items are linearly transformed to a 0-100 scale as follows: 0=0, 1=25, 2=50, 3=75, 4=100.</p> <p><u>Step 2: Calculate Scores</u></p> <p><u>Score by Dimensions:</u></p> <ul style="list-style-type: none"> • If more than 50% of the items in the scale are missing, the scale score should not be computed. • Mean score = Sum of the items over the number of items answered. • The General Health item is not included in the scale score, but is reported separately.
Interpretation and Analysis of Missing Data	<p>If more than 50% of the items in the scale are missing, the Scale Score should not be computed.</p> <p>If 50% or more items are completed: Impute the mean of the completed items in a scale.</p>

PedsQL™ 3.0 Arthritis Module

PARENT Report for Toddlers (ages 2-4)

The **Parent Report for Toddlers** (ages 2-4) of the **PedsQL™ 3.0 Arthritis Module** is composed of 14 items comprising 3 dimensions.

DESCRIPTION OF THE ARTHRITIS MODULE:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
Pain and Hurt	4	1-4	1-4	Higher scores indicate lower problems.
Daily Activities	5	1-5	1-5	
Treatment	5	1-5	1-5	

SCORING OF DIMENSIONS:

Item Scaling	5-point Likert scale from 0 (Never) to 4 (Almost always)
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed on a scale from 0 to 100.
Scoring Procedure	<p><u>Step 1: Transform Score</u></p> <p>Items are reversed scored and linearly transformed to a 0-100 scale as follows: 0=100, 1=75, 2=50, 3=25, 4=0.</p> <p><u>Step 2: Calculate Scores by Dimensions</u></p> <ul style="list-style-type: none"> • If more than 50% of the items in the scale are missing, the scale scores should not be computed • Mean score = Sum of the items over the number of items answered <p>There is no Total Score</p>
Interpretation and Analysis of Missing Data	<p>If more than 50% of the items in the scale are missing, the Scale Scores should not be computed.</p> <p>If 50% or more items are completed: Impute the mean of the completed items in a scale.</p>

PedsQL™ 3.0 Arthritis Module

CHILD and PARENT Reports for Young Children (ages 5-7)

The **Child and Parent Reports** of the **PedsQL™ 3.0 Arthritis Module** for **Young Children** (ages 5-7) are composed of 20 items comprising 5 dimensions.

DESCRIPTION OF THE ARTHRITIS MODULE:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
Pain and Hurt	4	1-4	1-4	Higher scores indicate lower problems.
Daily Activities	5	1-5	1-5	
Treatment	5	1-5	1-5	
Worry	3	1-3	1-3	
Communication	3	1-3	1-3	

SCORING OF DIMENSIONS:

Item Scaling	5-point Likert scale from 0 (Never) to 4 (Almost always) for the parent report 3-point scale: 0 (Not at all), 2 (Sometimes) and 4 (A lot) for the child report
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed on a scale from 0 to 100.
Scoring Procedure	<p><u>Step 1: Transform Score</u></p> <p>Items are reversed scored and linearly transformed to a 0-100 scale as follows: 0=100, 1=75, 2=50, 3=25, 4=0.</p> <p><u>Step 2: Calculate Scores by Dimensions</u></p> <ul style="list-style-type: none"> • If more than 50% of the items in the scale are missing, the scale scores should not be computed, • Mean score = Sum of the items over the number of items answered. <p>There is no Total Score</p>
Interpretation and Analysis of Missing Data	<p>If more than 50% of the items in the scale are missing, the Scale Scores should not be computed.</p> <p>If 50% or more items are completed: Impute the mean of the completed items in a scale.</p>

PedsQL™ 3.0 Arthritis Module

**CHILD and PARENT Reports for Children (ages 8-12), Teens
(ages 13-18)**

The **Child and Parent Reports** of the **PedsQL™ 3.0 Arthritis Module** for:

- Children (ages 8-12),
- And Teens (ages 13-18),

are composed of 22 items comprising 5 dimensions.

DESCRIPTION OF THE ARTHRITIS MODULE:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
Pain and Hurt	4	1-4	1-4	Higher scores indicate lower problems.
Daily Activities	5	1-5	1-5	
Treatment	7	1-7	1-7	
Worry	3	1-3	1-3	
Communication	3	1-3	1-3	

SCORING OF DIMENSIONS:

Item Scaling	5-point Likert scale from 0 (Never) to 4 (Almost always)
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed on a scale from 0 to 100.
Scoring Procedure	<p><u>Step 1: Transform Score</u></p> <p>Items are reversed scored and linearly transformed to a 0-100 scale as follows: 0=100, 1=75, 2=50, 3=25, 4=0.</p> <p><u>Step 2: Calculate Scores by Dimensions</u></p> <ul style="list-style-type: none"> • If more than 50% of the items in the scale are missing, the scale scores should not be computed, • Mean score = Sum of the items over the number of items answered. <p><u>There is no Total Score</u></p>
Interpretation and Analysis of missing data	<p>If more than 50% of the items in the scale are missing, the Scale Scores should not be computed.</p> <p>If 50% or more items are completed: Impute the mean of the completed items in a scale.</p>

PedsQL™ 3.0 Asthma Module

PARENT Report for Toddlers (ages 2-4)

The **Parent Report for Toddlers** (ages 2-4) of the **PedsQL™ 3.0 Asthma Module** is composed of 26 items comprising 4 dimensions.

DESCRIPTION OF THE ASTHMA MODULE:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
Asthma	11	1-11	1-11	Higher scores indicate lower problems.
Treatment	9	1-9	1-9	
Worry	3	1-3	1-3	
Communication	3	1-3	1-3	

SCORING OF DIMENSIONS:

Item Scaling	5-point Likert scale from 0 (Never) to 4 (Almost always)
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed on a scale from 0 to 100.
Scoring Procedure	<p><u>Step 1: Transform Score</u></p> <p>Items are reversed scored and linearly transformed to a 0-100 scale as follows: 0=100, 1=75, 2=50, 3=25, 4=0.</p> <p><u>Step 2: Calculate Scores by Dimensions</u></p> <ul style="list-style-type: none"> • If more than 50% of the items in the scale are missing, the scale scores should not be computed, • Mean score = Sum of the items over the number of items answered. <p>There is no Total Score</p>
Interpretation and Analysis of Missing Data	<p>If more than 50% of the items in the scale are missing, the Scale Scores should not be computed.</p> <p>If 50% or more items are completed: Impute the mean of the completed items in a scale.</p>

PedsQL[™] 3.0 Asthma Module

**CHILD and PARENT Reports for Young Children (ages 5-7),
Children (ages 8-12), Teens (ages 13-18)**

The **Child and Parent Reports** of the **PedsQL™ 3.0 Asthma Module** for:

- Young Children (ages 5-7),
- Children (ages 8-12),
- And Teens (ages 13-18),

are composed of 28 items comprising 4 dimensions.

DESCRIPTION OF THE ASTHMA MODULE:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
Asthma	11	1-11	1-11	Higher scores indicate lower problems.
Treatment	11	1-11	1-11	
Worry	3	1-3	1-3	
Communication	3	1-3	1-3	

SCORING OF DIMENSIONS:

Item Scaling	5-point Likert scale from 0 (Never) to 4 (Almost always) 3-point scale: 0 (Not at all), 2 (Sometimes) and 4 (A lot) for the child report for Young Children (ages 5-7)
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed on a scale from 0 to 100.
Scoring Procedure	<p><u>Step 1: Transform Score</u></p> <p>Items are reversed scored and linearly transformed to a 0-100 scale as follows: 0=100, 1=75, 2=50, 3=25, 4=0.</p> <p><u>Step 2: Calculate Scores by Dimensions</u></p> <ul style="list-style-type: none"> • If more than 50% of the items in the scale are missing, the scale scores should not be computed, • Mean score = Sum of the items over the number of items answered. <p>There is no Total Score</p>
Interpretation and Analysis of Missing Data	<p>If more than 50% of the items in the scale are missing, the Scale Scores should not be computed.</p> <p>If 50% or more items are completed: Impute the mean of the completed items in a scale.</p>

PedsQL[™] 3.0 SF22
Short Form Asthma Module

PARENT Report for Toddlers (ages 2-4)

The **Parent Report for Toddlers** (ages 2-4) of the **PedsQL™ 3.0 SF22 Asthma Module** is composed of 20 items comprising 2 dimensions.

DESCRIPTION OF THE SF22 ASTHMA MODULE:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
Asthma	11	1-11	1-11	Higher scores indicate lower problems.
Treatment	9	1-9	1-9	

SCORING OF DIMENSIONS:

Item Scaling	5-point Likert scale from 0 (Never) to 4 (Almost always)
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed on a scale from 0 to 100.
Scoring Procedure	<p><u>Step 1: Transform Score</u></p> <p>Items are reversed scored and linearly transformed to a 0-100 scale as follows: 0=100, 1=75, 2=50, 3=25, 4=0.</p> <p><u>Step 2: Calculate Scores by Dimensions</u></p> <ul style="list-style-type: none"> • If more than 50% of the items in the scale are missing, the scale scores should not be computed, • Mean score = Sum of the items over the number of items answered. <p>There is no Total Score</p>
Interpretation and Analysis of Missing Data	<p>If more than 50% of the items in the scale are missing, the Scale Scores should not be computed.</p> <p>If 50% or more items are completed: Impute the mean of the completed items in a scale.</p>

PedsQL™ 3.0 SF22
Short Form Asthma Module

**CHILD and PARENT Reports for Young Children (ages 5-7),
Children (ages 8-12), Teens (ages 13-18)**

The **Child and Parent Reports** of the **PedsQL™ 3.0 SF22 Asthma Module** for:

- Young Children (ages 5-7),
- Children (ages 8-12),
- And Teens (ages 13-18)

are composed of 22 items comprising 2 dimensions.

DESCRIPTION OF THE SF22 ASTHMA MODULE:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
Asthma	11	1-11	1-11	Higher scores indicate lower problems.
Treatment	11	1-11	1-11	

SCORING OF DIMENSIONS:

Item Scaling	5-point Likert scale from 0 (Never) to 4 (Almost always) 3-point scale: 0 (Not at all), 2 (Sometimes) and 4 (A lot) for the Child R Report for Young Children (ages 5-7)
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed on a scale from 0 to 100.
Scoring Procedure	<p><u>Step 1: Transform Score</u></p> <p>Items are reversed scored and linearly transformed to a 0-100 scale as follows: 0=100, 1=75, 2=50, 3=25, 4=0.</p> <p><u>Step 2: Calculate Scores by Dimensions</u></p> <ul style="list-style-type: none"> • If more than 50% of the items in the scale are missing, the scale scores should not be computed, • Mean score = Sum of the items over the number of items answered. <p>There is no Total Score</p>
Interpretation and Analysis of Missing Data	<p>If more than 50% of the items in the scale are missing, the Scale Scores should not be computed.</p> <p>If 50% or more items are completed: Impute the mean of the completed items in a scale.</p>

PedsQL[™] Brain Tumor Module

PARENT Report for Toddlers (ages 2-4)

The **Parent Report for Toddlers** (ages 2-4) of the **PedsQL™ Brain Tumor Module** is composed of 17 items comprising 5 dimensions.

DESCRIPTION OF THE BRAIN TUMOR MODULE:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
Pain and Hurt	3	1-3	1-3	Higher scores indicate lower problems.
Movement and Balance	3	1-3	1-3	
Procedural Anxiety	3	1-3	1-3	
Nausea	5	1-5	1-5	
Worry	3	1-3	1-3	

SCORING OF DIMENSIONS:

Item Scaling	5-point Likert scale from 0 (Never a problem) to 4 (Almost always a problem)
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed on a scale from 0 to 100.
Scoring Procedure	<p><u>Step 1: Transform Score</u></p> <p>Items are reversed scored and linearly transformed to a 0-100 scale as follows: 0=100, 1=75, 2=50, 3=25, 4=0.</p> <p><u>Step 2: Calculate Scores by Dimensions</u></p> <ul style="list-style-type: none"> • If more than 50% of the items in the scale are missing, the scale scores should not be computed, • Mean score = Sum of the items over the number of items answered. <p><u>There is no Total Score</u></p>
Interpretation and Analysis of Missing Data	<p>If more than 50% of the items in the scale are missing, the Scale Scores should not be computed.</p> <p>If 50% or more items are completed: Impute the mean of the completed items in a scale.</p>

PedsQL[™] Brain Tumor Module

CHILD and PARENT Reports for Young Children (ages 5-7)

The **Parent and Child Reports for Young Children** (ages 5-7) of the **PedsQL™ Brain Tumor Module** are composed of 23 items comprising 6 dimensions.

DESCRIPTION OF THE BRAIN TUMOR MODULE:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
Cognitive Problems	6	1-6	1-6	Higher scores indicate lower problems.
Pain and Hurt	3	1-3	1-3	
Movement and Balance	3	1-3	1-3	
Procedural Anxiety	3	1-3	1-3	
Nausea	5	1-5	1-5	
Worry	3	1-3	1-3	

SCORING OF DIMENSIONS:

Item Scaling	5-point Likert scale from 0 (Never) to 4 (Almost always) 3-point scale: 0 (Not at all), 2 (Sometimes) and 4 (A lot) for the Child Report for Young Children (ages 5-7)
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed on a scale from 0 to 100.
Scoring Procedure	<p><u>Step 1: Transform Score</u></p> <p>Items are reversed scored and linearly transformed to a 0-100 scale as follows: 0=100, 1=75, 2=50, 3=25, 4=0.</p> <p><u>Step 2: Calculate Scores by Dimensions</u></p> <ul style="list-style-type: none"> • If more than 50% of the items in the scale are missing, the scale scores should not be computed, • Mean score = Sum of the items over the number of items answered. <p>There is no Total Score</p>
Interpretation and Analysis of Missing Data	<p>If more than 50% of the items in the scale are missing, the Scale Scores should not be computed.</p> <p>If 50% or more items are completed: Impute the mean of the completed items in a scale.</p>

PedsQL[™] Brain Tumor Module

**CHILD and PARENT Reports for Children (ages 8-12), Teens
(ages 13-18)**

The **Parent and Child Reports for Children** (ages 8-12) and **Teens** (ages 13-18) of the **PedsQL™ Brain Tumor Module** are composed of 24 items comprising 6 dimensions.

DESCRIPTION OF THE BRAIN TUMOR MODULE:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
Cognitive Problems	7	1-7	1-7	Higher scores indicate lower problems
Pain and Hurt	3	1-3	1-3	
Movement and Balance	3	1-3	1-3	
Procedural Anxiety	3	1-3	1-3	
Nausea	5	1-5	1-5	
Worry	3	1-3	1-3	

SCORING OF DIMENSIONS:

Item Scaling	5-point Likert scale from 0 (Never a problem) to 4 (Almost always a problem)
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed on a scale from 0 to 100.
Scoring Procedure	<p><u>Step 1: Transform Score</u></p> <p>Items are reversed scored and linearly transformed to a 0-100 scale as follows: 0=100, 1=75, 2=50, 3=25, 4=0.</p> <p><u>Step 2: Calculate Scores by Dimensions</u></p> <ul style="list-style-type: none"> • If more than 50% of the items in the scale are missing, the scale scores should not be computed, • Mean score = Sum of the items over the number of items answered. <p><u>There is no Total Score</u></p>
Interpretation and Analysis of Missing Data	<p>If more than 50% of the items in the scale are missing, the Scale Scores should not be computed.</p> <p>If 50% or more items are completed: Impute the mean of the completed items in a scale.</p>

PedsQL™ 3.0 Cancer Module

PARENT Report for Toddlers (ages 2-4)

The Parent Report for Toddlers (ages 2-4) of the **PedsQL™ 3.0 Cancer Module** is composed of 25 items comprising 8 dimensions.

DESCRIPTION OF THE CANCER MODULE:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
Pain and Hurt	2	1-2	1-2	Higher scores indicate lower problems.
Nausea	5	1-5	1-5	
Procedural Anxiety	3	1-3	1-3	
Treatment Anxiety	3	1-3	1-3	
Worry	3	1-3	1-3	
Cognitive Problems	3	1-3	1-3	
Perceived Physical Appearance	3	1-3	1-3	
Communication	3	1-3	1-3	

SCORING OF DIMENSIONS:

Item Scaling	5-point Likert scale from 0 (Never) to 4 (Almost always)
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed to a 0 to 100 scale.
Scoring Procedure	<p><u>Step 1: Transform Score</u></p> <p>Items are reversed scored and linearly transformed to a 0-100 scale as follows: 0=100, 1=75, 2=50, 3=25, 4=0.</p> <p><u>Step 2: Calculate Scores by Dimensions</u></p> <ul style="list-style-type: none"> • If more than 50% of the items in the scale are missing, the scale scores should not be computed, • Mean score = Sum of the items over the number of items answered. <p><u>Total Score:</u> Sum of all the items over the number of items answered on all the Scales.</p>
Interpretation and Analysis of Missing Data	<p>If more than 50% of the items in the scale are missing, the Scale Scores should not be computed.</p> <p>If 50% or more items are completed: Impute the mean of the completed items in a scale.</p>

PedsQL™ 3.0 Cancer Module

CHILD and PARENT Reports for Young Children (ages 5-7)

The **Child and Parent Reports** of the **PedsQL™ 3.0 Cancer Module** for Young Children (ages 5-7), are composed of 26 items comprising 8 dimensions.

DESCRIPTION OF THE CANCER MODULE:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
Pain and Hurt	2	1-2	1-2	Higher scores indicate lower problems.
Nausea	5	1-5	1-5	
Procedural Anxiety	3	1-3	1-3	
Treatment Anxiety	3	1-3	1-3	
Worry	3	1-3	1-3	
Cognitive Problems	4	1-4	1-4	
Perceived Physical Appearance	3	1-3	1-3	
Communication	3	1-3	1-3	

SCORING OF DIMENSIONS:

Item Scaling	5-point Likert scale from 0 (Never) to 4 (Almost always) 3-point scale: 0 (Not at all), 2 (Sometimes) and 4 (A lot) for the Child Report for Young Children (ages 5-7)
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed on a scale from 0 to 100.
Scoring Procedure	<p><u>Step 1: Transform Score</u></p> <p>Items are reversed scored and linearly transformed to a 0-100 scale as follows: 0=100, 1=75, 2=50, 3=25, 4=0.</p> <p><u>Step 2: Calculate Scores by Dimensions</u></p> <ul style="list-style-type: none"> • If more than 50% of the items in the scale are missing, the scale scores should not be computed, • Mean score = Sum of the items over the number of items answered. <p><u>Total Score:</u> Sum of all the items over the number of items answered on all the Scales.</p>
Interpretation and Analysis of Missing Data	If more than 50% of the items in the scale are missing, the Scale Scores should not be computed. If 50% or more items are completed: Impute the mean of the completed items in a scale.

PedsQL[™] 3.0 Cancer Module

**CHILD and PARENT Reports for Children (ages 8-12), and
Teens (ages 13-18), and Young Adults Report (ages 18-25)**

The **Child and Parent Reports** of the **PedsQL™ 3.0 Cancer Module** for:

- Children (ages 8-12),
- Teens (ages 13-18)

and the Young Adults report (ages 18-25) are composed of 27 items comprising 8 dimensions.

DESCRIPTION OF THE CANCER MODULE:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
Pain and Hurt	2	1-2	1-2	Higher scores indicate lower problems.
Nausea	5	1-5	1-5	
Procedural Anxiety	3	1-3	1-3	
Treatment Anxiety	3	1-3	1-3	
Worry	3	1-3	1-3	
Cognitive Problems	5	1-5	1-5	
Perceived Physical Appearance	3	1-3	1-3	
Communication	3	1-3	1-3	

SCORING OF DIMENSIONS:

Item Scaling	5-point Likert scale from 0 (Never) to 4 (Almost always) 3-point scale: 0 (Not at all), 2 (Sometimes) and 4 (A lot) for the Child Report for Young Children (ages 5-7)
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed on a scale from 0 to 100.
Scoring Procedure	<p><u>Step 1: Transform Score</u></p> <p>Items are reversed scored and linearly transformed to a 0-100 scale as follows: 0=100, 1=75, 2=50, 3=25, 4=0.</p> <p><u>Step 2: Calculate Scores by Dimensions</u></p> <ul style="list-style-type: none"> • If more than 50% of the items in the scale are missing, the scale scores should not be computed, • Mean score = Sum of the items over the number of items answered. <p><u>Total Score:</u> Sum of all the items over the number of items answered on all the Scales.</p>
Interpretation and Analysis of Missing Data	If more than 50% of the items in the scale are missing, the Scale Scores should not be computed. If 50% or more items are completed: Impute the mean of the completed items in a scale.

PedsQL[™] 3.0 Cardiac Module

PARENT Report for Toddlers (ages 2-4)

The Parent Report for Toddlers (age 2-4) of the PedsQL™ 3.0 Cardiac Module is composed of 23 items comprising 6 dimensions.

DESCRIPTION OF THE CARDIAC MODULE:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
Heart problems and Treatment	7	1-7	1-7	Higher scores indicate lower problems.
Treatment II	3	1-3	1-3	
Perceived Physical Appearance	3	1-3	1-3	
Treatment Anxiety	4	1-4	1-4	
Cognitive Problems	3	1-3	1-3	
Communication	3	1-3	1-3	

SCORING OF DIMENSIONS:

Item Scaling	5-point Likert scale from 0 (Never) to 4 (Almost always)
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed to a 0 to 100 scale.
Scoring Procedure	<p><u>Step 1: Transform Score</u></p> <p>Items are reversed scored and linearly transformed to a 0-100 scale as follows: 0=100, 1=75, 2=50, 3=25, 4=0.</p> <p><u>Step 2: Calculate scores by Dimensions</u></p> <ul style="list-style-type: none"> • If more than 50% of the items in the scale are missing, the scale scores should not be computed, • Mean score = Sum of the items over the number of items answered. <p>There is no Total Score.</p>
Interpretation and Analysis of Missing Data	<p>If more than 50% of the items in the scale are missing, the Scale Scores should not be computed.</p> <p>If 50% or more items are completed: Impute the mean of the completed items in a scale.</p>

PedsQL[™] 3.0 Cardiac Module

CHILD and PARENT Reports for Young Children (ages 5-7)

The **Child and Parent Reports** of the **PedsQL™ 3.0 Cardiac Module** for Young Children (ages 5-7) are composed of 25 items comprising 6 dimensions.

DESCRIPTION OF THE CARDIAC MODULE:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
Heart Problems and Treatment	7	1-7	1-7	Higher scores indicate lower problems.
Treatment II	3	1-3	1-3	
Perceived Physical Appearance	3	1-3	1-3	
Treatment Anxiety	4	1-4	1-4	
Cognitive Problems	5	1-5	1-5	
Communication	3	1-3	1-3	

SCORING OF DIMENSIONS:

Item Scaling	5-point Likert scale from 0 (Never) to 4 (Almost always) 3-point scale: 0 (Not at all), 2 (Sometimes) and 4 (A lot) for the Child Report for Young Children (ages 5-7)
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed on a scale from 0 to 100.
Scoring Procedure	<p><u>Step 1: Transform Score</u></p> <p>Items are reversed scored and linearly transformed to a 0-100 scale as follows: 0=100, 1=75, 2=50, 3=25, 4=0</p> <p><u>Step 2: Calculate Scores by Dimensions</u></p> <ul style="list-style-type: none"> • If more than 50% of the items in the scale are missing, the scale scores should not be computed, • Mean score = Sum of the items over the number of items answered. <p>There is no Total Score</p>
Interpretation and Analysis of Missing Data	<p>If more than 50% of the items in the scale are missing, the Scale Scores should not be computed.</p> <p>If 50% or more items are completed: Impute the mean of the completed items in a scale.</p>

PedsQL[™] 3.0 Cardiac Module

**CHILD and PARENT Reports for Children (ages 8-12) and
Teens (ages 13-18)**

The **Child and Parent Reports** of the **PedsQL™ 3.0 Cardiac Module** for:

- Children (ages 8-12),
- And Teens (ages 13-18)

are composed of 27 items comprising 6 dimensions.

DESCRIPTION OF THE CARDIAC MODULE:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
Heart Problems and Treatment	7	1-7	1-7	Higher scores indicate lower problems.
Treatment II	5	1-5	1-5	
Perceived Physical Appearance	3	1-3	1-3	
Treatment Anxiety	4	1-4	1-4	
Cognitive Problems	5	1-5	1-5	
Communication	3	1-3	1-3	

SCORING OF DIMENSIONS:

Item Scaling	5-point Likert scale from 0 (Never) to 4 (Almost always)
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed on a scale from 0 to 100.
Scoring Procedure	<p><u>Step 1: Transform Score</u></p> <p>Items are reversed scored and linearly transformed to a 0-100 scale as follows: 0=100, 1=75, 2=50, 3=25, 4=0</p> <p><u>Step 2: Calculate Scores by Dimensions</u></p> <ul style="list-style-type: none"> • If more than 50% of the items in the scale are missing, the scale scores should not be computed, • Mean score = Sum of the items over the number of items answered. <p>There is no Total Score</p>
Interpretation and Analysis of Missing Data	<p>If more than 50% of the items in the scale are missing, the Scale Scores should not be computed.</p> <p>If 50% or more items are completed: Impute the mean of the completed items in a scale.</p>

PedsQL™ 3.0 Cerebral Palsy Module

PARENT Report for Toddlers (ages 2-4)

The Parent Report for Toddlers (ages 2-4) of the **PedsQL™ 3.0 Cerebral Palsy Module** is composed of 22 items comprising 5 dimensions.

DESCRIPTION OF THE CEREBRAL PALSY MODULE:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
Daily Activities	5	1-5	1-5	Higher scores indicate lower problems.
Movement and Balance	5	1-5	1-5	
Pain and Hurt	4	1-4	1-4	
Fatigue	4	1-4	1-4	
Eating Activities	4	1-4	1-4	

SCORING OF DIMENSIONS:

Item Scaling	5-point Likert scale from 0 (Never) to 4 (Almost always)
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed to a 0 to 100 scale.
Scoring Procedure	<p><u>Step 1: Transform Score</u></p> <p>Items are reversed scored and linearly transformed to a 0-100 scale as follows: 0=100, 1=75, 2=50, 3=25, 4=0</p> <p><u>Step 2: Calculate Scores by Dimensions</u></p> <ul style="list-style-type: none"> • If more than 50% of the items in the scale are missing, the scale scores should not be computed, • Mean score = Sum of the items over the number of items answered. <p>There is no Total score</p>
Interpretation and Analysis of Missing Data	<p>If more than 50% of the items in the scale are missing, the Scale Scores should not be computed.</p> <p>If 50% or more items are completed: Impute the mean of the completed items in a scale.</p>

PedsQL[™] 3.0 Cerebral Palsy Module

**CHILD and PARENT Reports for Young Children (ages 5-7),
Children (ages 8-12), Teens (ages 13-18)**

The **Child and Parent Reports** of the **PedsQL™ 3.0 Cerebral Palsy Module** for:

- Young Children (ages 5-7),
- Children (ages 8-12),
- And Teens (ages 13-18)

are composed of 35 items comprising 7 dimensions.

DESCRIPTION OF THE CEREBRAL PALSY MODULE:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
Daily Activities	9	1-9	1-9	Higher scores indicate lower problems.
School Activities	4	1-4	1-4	
Movement and Balance	5	1-5	1-5	
Pain and Hurt	4	1-4	1-4	
Fatigue	4	1-4	1-4	
Eating Activities	5	1-5	1-5	
Speech and Communication	4	1-4	1-4	

SCORING OF DIMENSIONS:

Item Scaling	5-point Likert scale from 0 (Never) to 4 (Almost always) 3-point scale: 0 (Not at all), 2 (Sometimes) and 4 (A lot) for the Child Report for Young Children (ages 5-7)
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed on a scale from 0 to 100.
Scoring Procedure	<p><u>Step 1: Transform Score</u> Items are reversed scored and linearly transformed to a 0-100 scale as follows: 0=100, 1=75, 2=50, 3=25, 4=0.</p> <p><u>Step 2: Calculate Scores by Dimensions</u></p> <ul style="list-style-type: none"> • If more than 50% of the items in the scale are missing, the scale scores should not be computed, • Mean score = Sum of the items over the number of items answered. <p>There is no Total score</p>
Interpretation and Analysis of Missing Data	If more than 50% of the items in the scale are missing, the Scale Scores should not be computed. If 50% or more items are completed: Impute the mean of the completed items in a scale.

PedsQL™ Cognitive Functioning Scale

The **PedsQL™ Cognitive Functioning Scale** is composed of 6 items comprising 1 dimension.

DESCRIPTION OF THE COGNITIVE FUNTIONING SCALE:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
Cognitive Functioning	6	1-6	1-6	Higher scores indicate lower problems

SCORING OF DIMENSIONS:

Item Scaling	5-point Likert scale from 0 (Never) to 4 (Almost always) 3-point scale: 0 (Not at all), 2 (Sometimes) and 4 (A lot) for the Child Report for Young Children (ages 5-7)
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed on a scale from 0 to 100.
Scoring Procedure	<p><u>Step 1: Transform Score</u></p> <p>Items are reversed scored and linearly transformed to a 0-100 scale as follows: 0=100, 1=75, 2=50, 3=25, 4=0.</p> <p><u>Step 2: Calculate Total Score</u></p> <p>Sum of all the items over the number of items answered</p>
Interpretation and Analysis of Missing Data	<p>If more than 50% of the items in the scale are missing, the Scale Score should not be computed.</p> <p>If 50% or more items are completed: Impute the mean of the completed items in a scale.</p>

PedsQL[™] 3.0 Diabetes Module

The **PedsQL™ 3.0 Diabetes Module** is composed of 28 items comprising 5 dimensions.

DESCRIPTION OF THE DIABETES MODULE:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
Diabetes	11	1-11	1-11	Higher scores indicate lower problems.
Treatment I	4	1-4	1-4	
Treatment II	7	1-7	1-7	
Worry	3	1-3	1-3	
Communication	3	1-3	1-3	

SCORING OF DIMENSIONS:

Item Scaling	5-point scale from 0 (Never) to 4 (Almost always) 3-point scale: 0 (Not at all), 2 (Sometimes) and 4 (A lot) for the Child Report for Young Children (ages 5-7)
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed on a scale from 0 to 100.
Scoring Procedure	<p><u>Step 1: Transform Score</u></p> <p>Items are reversed scored and linearly transformed to a 0-100 scale as follows: 0=100, 1=75, 2=50, 3=25, 4=0.</p> <p><u>Step 2: Calculate scores by Dimension</u></p> <ul style="list-style-type: none"> • If more than 50% of the items in the scale are missing, the scale scores should not be computed, • Mean score = Sum of the items over the number of items answered. <p><u>Total Score:</u> Sum of all the items over the number of items answered on all the Scales.</p>
Interpretation and Analysis of Missing Data	<p>If more than 50% of the items in the scale are missing, the Scale Scores should not be computed.</p> <p>If 50% or more items are completed: Impute the mean of the completed items in a scale.</p>

PedsQL[™] 3.2 Diabetes Module

The **PedsQL™ 3.2 Diabetes Module** is composed of 33 items comprising 5 dimensions for ages 13-45 years. For ages 2-12 years, the **PedsQL™ 3.2 Diabetes Module** is composed of 32 items comprising 5 dimensions (one less item for the Worry Scale).

DESCRIPTION OF THE DIABETES MODULE:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
Diabetes	15	1-15	1-15	Higher scores indicate lower problems.
Treatment I	5	1-5	1-5	
Treatment II	6	1-6	1-6	
Worry	2 (3 for teens and adults)	1-2 or 1-3	1-2 or 1-3	
Communication	4	1-4	1-4	

SCORING OF DIMENSIONS:

Item Scaling	5-point scale from 0 (Never) to 4 (Almost always) 3-point scale: 0 (Not at all), 2 (Sometimes) and 4 (A lot) for the Child Report for Young Children (ages 5-7)
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed on a scale from 0 to 100.
Scoring Procedure	<p><u>Step 1: Transform Score</u></p> <p>Items are reversed scored and linearly transformed to a 0-100 scale as follows: 0=100, 1=75, 2=50, 3=25, 4=0.</p> <p><u>Step 2: Calculate scores by Dimension</u></p> <ul style="list-style-type: none"> • If more than 50% of the items in the scale are missing, the scale scores should not be computed, • Mean score = Sum of the items over the number of items answered. <p><u>Total Score:</u> Sum of all the items over the number of items answered on all the Scales.</p>
Interpretation and Analysis of Missing Data	<p>If more than 50% of the items in the scale are missing, the Scale Scores should not be computed.</p> <p>If 50% or more items are completed: Impute the mean of the completed items in a scale.</p>

PedsQL™ 3.0 Duchenne Muscular Dystrophy Module

PARENT Report for Young Children (ages 5-7)

The **Parent Report** of the **PedsQL™ 3.0 Duchenne Muscular Dystrophy Module** for Young Children (ages 5-7) is composed of 18 items comprising 4 dimensions.

DESCRIPTION OF THE DUCHENNE MUSCULAR DYSTROPHY MODULE:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
Daily Activities	5	1-5	1-5	Higher scores indicate lower problems.
Treatment	4	1-4	1-4	
Worry	6	1-6	1-6	
Communication	3	1-3	1-3	

SCORING OF DIMENSIONS:

Item Scaling	5-point Likert scale from 0 (Never) to 4 (Almost always)
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed on a scale from 0 to 100.
Scoring Procedure	<p><u>Step 1: Transform Score</u></p> <p>Items are reversed scored and linearly transformed to a 0-100 scale as follows: 0=100, 1=75, 2=50, 3=25, 4=0</p> <p><u>Step 2: Calculate Scores by Dimensions</u></p> <ul style="list-style-type: none"> If more than 50% of the items in the scale are missing, the scale scores should not be computed, Mean score = Sum of the items over the number of items answered. <p><u>No Total Score</u></p>
Interpretation and Analysis of Missing Data	<p>If more than 50% of the items in the scale are missing, the Scale Scores should not be computed.</p> <p>If 50% or more items are completed: Impute the mean of the completed items in a scale.</p>

PedsQL[™] 3.0 Duchenne Muscular Dystrophy Module

**CHILD and PARENT Reports for Children (ages 8-12) and
Teens (ages 13-18)**

The **Child and Parent Reports** of the **PedsQL™ 3.0 Duchenne Muscular Dystrophy Module** for:

- Children (ages 8-12),
- And Teens (ages 13-18)

are composed of 18 items comprising 4 dimensions.

DESCRIPTION OF THE DUCHENNE MUSCULAR DYSTROPHY MODULE:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
Daily Activities	5	1-5	1-5	Higher scores indicate lower problems.
Treatment	4	1-4	1-4	
Worry	6	1-6	1-6	
Communication	3	1-3	1-3	

SCORING OF DIMENSIONS:

Item Scaling	5-point Likert scale from 0 (Never) to 4 (Almost always)
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed on a scale from 0 to 100.
Scoring Procedure	<p><u>Step 1: Transform Score</u></p> <p>Items are reversed scored and linearly transformed to a 0-100 scale as follows: 0=100, 1=75, 2=50, 3=25, 4=0</p> <p><u>Step 2: Calculate Scores by Dimensions</u></p> <ul style="list-style-type: none"> • If more than 50% of the items in the scale are missing, the scale scores should not be computed, • Mean score = Sum of the items over the number of items answered. <p><u>No Total Score</u></p>
Interpretation and Analysis of Missing Data	<p>If more than 50% of the items in the scale are missing, the Scale Scores should not be computed.</p> <p>If 50% or more items are completed: Impute the mean of the completed items in a scale.</p>

PedsQL™ 3.0 Eosinophilic Esophagitis Module

PARENT Report for Toddlers (ages 2-4)

The Parent Report for Toddlers (age 2-4) of the PedsQL™ 3.0 Eosinophilic Esophagitis Module is composed of 21 items comprising 6 dimensions.

DESCRIPTION OF THE EOSINOPHILIC ESOPHAGITIS MODULE:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
Symptoms I	3	1-3	1-3	Higher scores indicate lower problems.
Symptoms II	4	1-4	1-4	
Treatment	4	1-4	1-4	
Worry	4	1-4	1-4	
Food and Eating	3	1-3	1-3	
Food Feelings	3	1-3	1-3	

Note: Feeding Tube Scale not scored.

SCORING OF DIMENSIONS:

Item Scaling	5-point Likert scale from 0 (Never) to 4 (Almost always)
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed to a 0 to 100 scale.
Scoring Procedure	<p><u>Step 1: Transform Score</u></p> <p>Items are reversed scored and linearly transformed to a 0-100 scale as follows: 0=100, 1=75, 2=50, 3=25, 4=0.</p> <p><u>Step 2: Calculate scores by Dimensions</u></p> <ul style="list-style-type: none"> • If more than 50% of the items in the scale are missing, the scale scores should not be computed, • Mean score = Sum of the items over the number of items answered. <p><u>Symptoms Total Scale Score</u> = Sum of the items over the number of items answered in the Symptoms I and Symptoms II Scales.</p> <p><u>Total Score:</u> Sum of all the items over the number of items answered on all the Scales except Feeding Tube Scale.</p>
Interpretation and Analysis of Missing Data	<p>If more than 50% of the items in the scale are missing, the Scale Scores should not be computed.</p> <p>If 50% or more items are completed: Impute the mean of the completed items in a scale.</p>

REFERENCES:

Franciosi, J.P., Hommel, K.A., Bendo, C.B., King, E.C., Collins, M.H., Eby, M.D., Marsolo, K., Abonia, J.P., von Tiehl, K.F., Putnam, P.E., Greenler, A.J., Greenberg, A.B., Bryson,

R.A., Davis, C.M., Olive, A.P., Gupta, S.K., Erwin, E.A., Klinnert, M.D., Spergel, J.M., Denham, J.M., Furuta, G.T., Rothenberg, M.E., & Varni, J.W. (2012). PedsQL™ Eosinophilic Esophagitis Module: Feasibility, reliability and validity. Manuscript submitted for publication.

Franciosi, J.P., Hommel, K.A., Greenberg, A.B., Debrosse, C.W., Greenler, A.J., Abonia, J.P., Rothenberg, M.E., & Varni, J.W. (2012). Development of the Pediatric Quality of Life Inventory™ Eosinophilic Esophagitis Module items: Qualitative methods. Manuscript submitted for publication.

Franciosi J.P., Hommel, K.A., Debrosse, C.W., Greenberg, A.B., Greenler, A.J., Abonia, J.P., Rothenberg, M.E., & Varni, J.W. (2012). Quality of life in paediatric eosinophilic oesophagitis: What is important to patients? *Child: Care, Health and Development*, 38, 477–483.

PedsQL[™] 3.0 Eosinophilic Esophagitis Module

CHILD and PARENT Reports for Young Children (ages 5-7)

The **Child and Parent Reports** of the **PedsQL™ 3.0 Eosinophilic Esophagitis Module** for Young Children (ages 5-7) are composed of 26 items comprising 7 dimensions.

DESCRIPTION OF THE EOSINOPHILIC ESOPHAGITIS MODULE:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
Symptoms I	3	1-3	1-3	Higher scores indicate lower problems.
Symptoms II*	4	1-4	1-4	
Treatment	4	1-4	1-4	
Worry	4	1-4	1-4	
Communication	4	1-4	1-4	
Food and Eating	4	1-4	1-4	
Food Feelings	3	1-3	1-3	

*Symptoms II Scale not scored for child self-report ages 5-7 years.
 Note: Feeding Tube Scale not scored.

SCORING OF DIMENSIONS:

Item Scaling	5-point Likert scale from 0 (Never) to 4 (Almost always) 3-point scale: 0 (Not at all), 2 (Sometimes) and 4 (A lot) for the Child Report for Young Children (ages 5-7)
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed on a scale from 0 to 100.
Scoring Procedure	<p><u>Step 1: Transform Score</u></p> <p>Items are reversed scored and linearly transformed to a 0-100 scale as follows: 0=100, 1=75, 2=50, 3=25, 4=0</p> <p><u>Step 2: Calculate Scores by Dimensions</u></p> <ul style="list-style-type: none"> • If more than 50% of the items in the scale are missing, the scale scores should not be computed, • Mean score = Sum of the items over the number of items answered. <p><u>Symptoms Total Scale Score</u> = Sum of the items over the number of items answered in the Symptoms I and Symptoms II Scales. No Symptoms Total Scale Score for child self-report ages 5-7 years.</p> <p><u>Total Score:</u> Sum of all the items over the number of items answered on all the Scales for parent report except Feeding Tube Scale. For child self-report ages 5-7, sum all of the items over the number of items answered on all Scales except for Symptoms II and Feeding Tube Scale.</p>

Interpretation and Analysis of Missing Data	If more than 50% of the items in the scale are missing, the Scale Scores should not be computed. If 50% or more items are completed: Impute the mean of the completed items in a scale.
--	--

REFERENCES:

Franciosi, J.P., Hommel, K.A., Bendo, C.B., King, E.C., Collins, M.H., Eby, M.D., Marsolo, K., Abonia, J.P., von Tiehl, K.F., Putnam, P.E., Greenler, A.J., Greenberg, A.B., Bryson, R.A., Davis, C.M., Olive, A.P., Gupta, S.K., Erwin, E.A., Klinnert, M.D., Spergel, J.M., Denham, J.M., Furuta, G.T., Rothenberg, M.E., & Varni, J.W. (2012). PedsQL™ Eosinophilic Esophagitis Module: Feasibility, reliability and validity. Manuscript submitted for publication.

Franciosi, J.P., Hommel, K.A., Greenberg, A.B., Debrosse, C.W., Greenler, A.J., Abonia, J.P., Rothenberg, M.E., & Varni, J.W. (2012). Development of the Pediatric Quality of Life Inventory™ Eosinophilic Esophagitis Module items: Qualitative methods. Manuscript submitted for publication.

Franciosi J.P., Hommel, K.A., Debrosse, C.W., Greenberg, A.B., Greenler, A.J., Abonia, J.P., Rothenberg, M.E., & Varni, J.W. (2012). Quality of life in paediatric eosinophilic oesophagitis: What is important to patients? *Child: Care, Health and Development*, 38, 477–483.

PedsQL[™] 3.0 Eosinophilic Esophagitis Module

**CHILD and PARENT Reports for Children (ages 8-12) and
Teens (ages 13-18)**

The **Child and Parent Reports** of the **PedsQL™ 3.0 Eosinophilic Esophagitis Module** for:
 - Children (ages 8-12),
 - And Teens (ages 13-18)
 are composed of 33 items comprising 7 dimensions.

DESCRIPTION OF THE EOSINOPHILIC ESOPHAGITIS MODULE:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
Symptoms I	6	1-6	1-6	Higher scores indicate lower problems.
Symptoms II	4	1-4	1-4	
Treatment	5	1-5	1-5	
Worry	6	1-6	1-6	
Communication	5	1-5	1-5	
Food and Eating	4	1-4	1-4	
Food Feelings	3	1-3	1-3	

Note: Feeding Tube Scale not scored.

SCORING OF DIMENSIONS:

Item Scaling	5-point Likert scale from 0 (Never) to 4 (Almost always)
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed on a scale from 0 to 100.
Scoring Procedure	<p><u>Step 1: Transform Score</u></p> <p>Items are reversed scored and linearly transformed to a 0-100 scale as follows: 0=100, 1=75, 2=50, 3=25, 4=0</p> <p><u>Step 2: Calculate Scores by Dimensions</u></p> <ul style="list-style-type: none"> • If more than 50% of the items in the scale are missing, the scale scores should not be computed, • Mean score = Sum of the items over the number of items answered. <p><u>Symptoms Total Scale Score</u> = Sum of the items over the number of items answered in the Symptoms I and Symptoms II Scales.</p> <p><u>Total Score:</u> Sum of all the items over the number of items answered on all the Scales except Feeding Tube Scale.</p>
Interpretation and Analysis of Missing Data	<p>If more than 50% of the items in the scale are missing, the Scale Scores should not be computed.</p> <p>If 50% or more items are completed: Impute the mean of the completed items in a scale.</p>

REFERENCES:

Franciosi, J.P., Hommel, K.A., Bendo, C.B., King, E.C., Collins, M.H., Eby, M.D., Marsolo, K., Abonia, J.P., von Tiehl, K.F., Putnam, P.E., Greenler, A.J., Greenberg, A.B., Bryson, R.A., Davis, C.M., Olive, A.P., Gupta, S.K., Erwin, E.A., Klinnert, M.D., Spergel, J.M., Denham, J.M., Furuta, G.T., Rothenberg, M.E., & Varni, J.W. (2012). PedsQL™ Eosinophilic Esophagitis Module: Feasibility, reliability and validity. Manuscript submitted for publication.

Franciosi, J.P., Hommel, K.A., Greenberg, A.B., Debrosse, C.W., Greenler, A.J., Abonia, J.P., Rothenberg, M.E., & Varni, J.W. (2012). Development of the Pediatric Quality of Life Inventory™ Eosinophilic Esophagitis Module items: Qualitative methods. Manuscript submitted for publication.

Franciosi J.P., Hommel, K.A., Debrosse, C.W., Greenberg, A.B., Greenler, A.J., Abonia, J.P., Rothenberg, M.E., & Varni, J.W. (2012). Quality of life in paediatric eosinophilic oesophagitis: What is important to patients? *Child: Care, Health and Development*, 38, 477–483.

PedsQL™ 3.0 End Stage Renal Disease Module

PARENT Report for Toddlers (ages 2-4)

The Parent Report for Toddlers (ages 2-4) of the **PedsQL™ 3.0 End Stage Renal Disease Module** is composed of 13 items comprising 4 dimensions.

DESCRIPTION OF THE END STAGE RENAL DISEASE MODULE:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
General Fatigue	3	1-3	1-3	Higher scores indicate lower problems.
About My Kidney Disease	4	1-4	1-4	
Treatment Problems	3	1-3	1-3	
Worry	3	1-3	1-3	

SCORING OF DIMENSIONS:

Item Scaling	5-point Likert scale from 0 (Never) to 4 (Almost always)
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed to a 0 to 100 scale.
Scoring Procedure	<p><u>Step 1: Transform Score</u></p> <p>Items are reversed scored and linearly transformed to a 0-100 scale as follows: 0=100, 1=75, 2=50, 3=25, 4=0.</p> <p><u>Step 2: Calculate Scores by Dimensions</u></p> <ul style="list-style-type: none"> • If more than 50% of the items in the scale are missing, the scale scores should not be computed, • Mean score = Sum of the items over the number of items answered. <p><u>Total Score:</u> Sum of all the items over the number of items answered on all the Scales.</p>
Interpretation and Analysis of Missing Data	<p>If more than 50% of the items in the scale are missing, the Scale Scores should not be computed.</p> <p>If 50% or more items are completed: Impute the mean of the completed items in a scale.</p>

PedsQL™ 3.0 End Stage Renal Disease Module

**CHILD and PARENT Reports for Young Children (ages 5-7),
Children (ages 8-12), Teens (ages 13-18)**

The **Child and Parent Reports** of the **PedsQL™ 3.0 End Stage Renal Disease Module** for:

- Young Children (ages 5-7),
 - Children (ages 8-12),
 - And Teens ages (13-18)
- are composed of 34 items comprising 7 dimensions.

DESCRIPTION OF THE END STAGE RENAL DISEASE MODULE:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
General Fatigue	4	1-4	1-4	Higher scores indicate lower problems.
About My Kidney Disease	5	1-5	1-5	
Treatment Problems	4	1-4	1-4	
Family and Peer Interaction	3	1-3	1-3	
Worry	10	1-10	1-10	
Perceived Physical Appearance	3	1-3	1-3	
Communication	5	1-5	1-5	

SCORING OF DIMENSIONS:

Item Scaling	5-point Likert scale from 0 (Never) to 4 (Almost always) 3-point scale: 0 (Not at all), 2 (Sometimes) and 4 (A lot) for the Young Children (ages 5-7) child report
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed on a scale from 0 to 100.
Scoring Procedure	<p><u>Step 1: Transform Score</u></p> <p>Items are reversed scored and linearly transformed to a 0-100 scale as follows: 0=100, 1=75, 2=50, 3=25, 4=0.</p> <p><u>Step 2: Calculate Scores</u></p> <p><u>Score by Dimensions:</u></p> <ul style="list-style-type: none"> • If more than 50% of the items in the scale are missing, the scale scores should not be computed, • Mean score = Sum of the items over the number of items answered. <p><u>Total Score:</u> Sum of all the items over the number of items answered on all the Scales.</p>
Interpretation and Analysis of Missing Data	<p>If more than 50% of the items in the scale are missing, the Scale Scores should not be computed.</p> <p>If 50% or more items are completed: Impute the mean of the completed items in a scale.</p>

PedsQL[™] 2.0 Family Impact Module

PARENT Report

The Parent report of the **PedsQL™ 2.0 Family impact Module** is composed of 36 items comprising 8 dimensions.

DESCRIPTION OF THE FAMILY IMPACT MODULE:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
Physical Functioning	6	1-6	1-6	Higher scores indicate better functioning.
Emotional Functioning	5	1-5	1-5	
Social Functioning	4	1-4	1-4	
Cognitive Functioning	5	1-5	1-5	
Communication	3	1-3	1-3	
Worry	5	1-5	1-5	
Daily Activities	3	1-3	1-3	
Family Relationships	5	1-5	1-5	

SCORING OF DIMENSIONS:

Item Scaling	5-point Likert scale from 0 (Never) to 4 (Almost always)
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed to a 0 to 100 scale.
Scoring Procedure	<p><u>Step 1: Transform Score</u></p> <p>Items are reversed scored and linearly transformed to a 0-100 scale as follows: 0=100, 1=75, 2=50, 3=25, 4=0</p> <p><u>Step 2: Calculate Scores by Dimensions</u></p> <ul style="list-style-type: none"> • If more than 50% of the items in the scale are missing, the scale scores should not be computed, • Mean score = Sum of the items over the number of items answered. <p><u>Step 3: Total Scores</u></p> <ul style="list-style-type: none"> • <u>The Total Score</u> is the sum of all 36 items divided by the number of items answered • <u>The Parent HRQL Summary Score</u> (20 items) is computed as the sum of the items divided by the number of items answered in the Physical, Emotional, Social, and Cognitive Functioning scales. • <u>The Family Functioning Summary Score</u> (8 items) is computed as the sum of the items divided by the number of items answered in the Daily Activities and family Relationships scales.

Interpretation and Analysis of Missing Data	If more than 50% of the items in the scale are missing, the Scale Scores should not be computed. If 50% or more items are completed: Impute the mean of the completed items in a scale.
--	--

PedsQL[™] Gastrointestinal Symptom Scale (GSS)

PARENT Proxy Report

The PARENT proxy report **PedsQL™ Gastrointestinal Symptom Scale** is composed of 9 items comprising 1 dimension.

DESCRIPTION OF THE GASTROINTESTINAL SYMPTOM SCALE:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
Gastrointestinal Symptoms	9	1-9	1-9	Higher scores indicate lower problems.

SCORING OF DIMENSIONS:

Item Scaling	5-point Likert scale from 0 (Never) to 4 (Almost always)
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed on a scale from 0 to 100.
Scoring Procedure	<p><u>Step 1: Transform Score</u></p> <p>Items are reversed scored and linearly transformed to a 0-100 scale as follows: 0=100, 1=75, 2=50, 3=25, 4=0.</p> <p><u>Step 2: Calculate Total Score</u></p> <p>Sum of all the items over the number of items answered on all the Scales.</p>
Interpretation and Analysis of Missing Data	<p>If more than 50% of the items in the scale are missing, the Scale Scores should not be computed.</p> <p>If 50% or more items are completed: Impute the mean of the completed items in a scale.</p>

PedsQL[™] Healthcare Satisfaction Generic Module

The **PedsQL™ Healthcare Satisfaction Generic Module** is composed of 24 items comprising 6 dimensions.

DESCRIPTION OF THE MODULE:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
Information	5	1-5	No	Higher scores indicate higher satisfaction.
Inclusion of Family	4	1-4		
Communication	5	1-5		
Technical Skills	3	1-3		
Emotional Needs	4	1-4		
Overall Satisfaction	3	1-3		

SCORING OF DIMENSIONS:

Item Scaling	5-point Likert scale: 0 (Never) to 4 (Always) and Not Applicable
Weighting of items	No
Extension of the Scoring Scale	Scores are transformed on a scale from 0 to 100.
Scoring Procedure	<p><u>Step 1: Transform Score</u></p> <p>Items are scored and linearly transformed to a 0-100 scale as follows: 0=0, 1=25, 2=50, 3=75, 4=100</p> <p><u>Step 2: Calculate Scores by Dimensions</u></p> <ul style="list-style-type: none"> • If 50% or more items are completed, the Mean score = Sum of the items divided by the number of items answered in the scale, • If more than 50% of the items in the scale are missing, the scale scores should not be computed. <p><u>Total Score:</u> Mean score = Sum of all the items in all scales divided by the number of items answered on all the scales.</p>
Interpretation and Analysis of Missing Data	<p>If more than 50% of the items in the scale are missing, the Scale Scores should not be computed.</p> <p>If 50% or more items are completed: Impute the mean of the completed items in the scale.</p>

PedsQL[™] Healthcare Satisfaction Hematology / Oncology Module

The **PedsQL™ Healthcare Satisfaction Hematology / Oncology Module** is composed of 25 items comprising 6 dimensions.

DESCRIPTION OF THE MODULE:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
General Satisfaction	3	1-3	No	Higher scores indicate higher satisfaction.
Information	5	1-5		
Inclusion of Family	4	1-4		
Communication	5	1-5		
Technical Skills	4	1-4		
Emotional Needs	4	1-4		

SCORING OF DIMENSIONS:

Item Scaling	5-point Likert scale: 1 (Very dissatisfied) to 5 (Very satisfied)
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed on a scale from 0 to 100
Scoring Procedure	<p><u>Step 1: Transform Score</u></p> <p>Items are scored and linearly transformed to a 0-100 scale as follows: 1=0, 2=25, 3=50, 4=75, 5=100.</p> <p><u>Step 2: Calculate Scores by Dimensions</u></p> <ul style="list-style-type: none"> • If 50% or more items are completed, the Mean score = Sum of the items divided by the number of items answered in the scale, • If more than 50% of the items in the scale are missing, the scale scores should not be computed. <p><u>Total Score:</u> Mean score = Sum of all the items in all scales divided by the number of items answered on all the scales.</p>
Interpretation and Analysis of Missing Data	<p>If more than 50% of the items in the scale are missing, the Scale Scores should not be computed.</p> <p>If 50% or more items are completed: Impute the mean of the completed items in the scale.</p>

PedsQL[™] Multidimensional Fatigue Scale

The **PedsQL™ Multidimensional Fatigue Scale** is composed of 18 items comprising 3 dimensions.

DESCRIPTION OF THE MULTIDIMENSIONAL FATIGUE MODULE:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
General Fatigue	6	1-6	1-6	Higher scores indicate lower problems.
Sleep/Rest Fatigue	6	1-6	1-6	
Cognitive Fatigue	6	1-6	1-6	

SCORING OF DIMENSIONS:

Item Scaling	5-point Likert scale from 0 (Never) to 4 (Almost always) 3-point scale: 0 (Not at all), 2 (Sometimes) and 4 (A lot) for the Child Report for Young Children (ages 5-7)
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed on a scale from 0 to 100.
Scoring Procedure	<p><u>Step 1: Transform Score</u></p> <p>Items are reversed scored and linearly transformed to a 0-100 scale as follows: 0=100, 1=75, 2=50, 3=25, 4=0.</p> <p><u>Step 2: Calculate Scores by Dimensions</u></p> <ul style="list-style-type: none"> • If more than 50% of the items in the scale are missing, the scale scores should not be computed, • Mean score= Sum of the items over the number of items answered. <p><u>Total Score:</u> Sum of all the items over the number of items answered on all the Scales</p>
Interpretation and Analysis of Missing Data	<p>If more than 50% of the items in the scale are missing, the Scale Scores should not be computed.</p> <p>If 50% or more items are completed: Impute the mean of the completed items in a scale.</p>

PedsQL[™] 1.0 Neurofibromatosis Type 1 Module

ADULT Report (≥ 26)

The **Adult Report** of the **PedsQL™ 1.0 Neurofibromatosis Type 1 Module** for:
 - Adults (ages ≥ 26)
 are composed of 70 items comprising 16 dimensions.

DESCRIPTION OF THE CEREBRAL PALSY MODULE:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
Physical Functioning	7	1-7	1-7	Higher scores indicate lower problems.
Emotional Functioning	5	1-5	1-5	
Social Functioning	2	1-2	1-2	
Cognitive Functioning	5	1-5	1-5	
Communication	3	1-3	1-3	
Worry	6	1-6	1-6	
Perceived Physical Appearance	3	1-3	1-3	
Pain and Hurt	3	1-3	1-3	
Paresthesias	2	1-2	1-2	
Skin Irritation	5	1-5	1-5	
Sensation	4	1-4	1-4	
Movement and Balance	4	1-4	1-4	
Daily Activities	12	1-12	1-12	
Feeling	3	1-3	1-3	
Treatment Anxiety	3	1-3	1-3	
Sexual Functioning	3	1-3	1-3	

SCORING OF DIMENSIONS:

Item Scaling	5-point Likert scale from 0 (Never) to 4 (Almost always)
Weighting of Items	No
Extension of the	Scores are transformed on a scale from 0 to 100.

Scoring Scale	
Scoring Procedure	<p><u>Step 1: Transform Score</u> Items are reversed scored and linearly transformed to a 0-100 scale as follows: 0=100, 1=75, 2=50, 3=25, 4=0.</p> <p><u>Step 2: Calculate Scores by Dimensions</u></p> <ul style="list-style-type: none"> • If more than 50% of the items in the scale are missing, the scale scores should not be computed, • Mean score = Sum of the items over the number of items answered. <p><u>Total Score:</u> Sum of all the items over the number of items answered on all the Scales.</p>
Interpretation and Analysis of Missing Data	<p>If more than 50% of the items in the scale are missing, the Scale Scores should not be computed.</p> <p>If 50% or more items are completed: Impute the mean of the completed items in a scale.</p>

PedsQL[™] 3.0 Neuromuscular Module

Child and Parent Reports of the **PedsQL™ 3.0 Neuromuscular Module** for:

- Children (ages 8-12),
- And Teens (ages 13-18),

are composed of 25 items comprising 3 dimensions.

- **Young Child Report** (ages 5-7) only consists of the 17-item About My Neuromuscular Disease Scale, **Parent Report** for ages 2-18 includes all 3 dimensions and all 25 items,

DESCRIPTION OF THE NEUROMUSCULAR MODULE:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
About My Neuromuscular Disease	17	1-17	1-17	Higher scores indicate lower problems.
Communication	3	1-3	1-3	
About Our Family Resources	5	1-5	1-5	

SCORING OF DIMENSIONS:

Item Scaling	5-point Likert scale from 0 (Never) to 4 (Almost always)
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed on a scale from 0 to 100.
Scoring Procedure	<p><u>Step 1: Transform Score</u></p> <p>Items are reversed scored and linearly transformed to a 0-100 scale as follows: 0=100, 1=75, 2=50, 3=25, 4=0.</p> <p><u>Step 2: Calculate Scores by Dimensions</u></p> <ul style="list-style-type: none"> • If more than 50% of the items in the scale are missing, the scale scores should not be computed, • Mean score = Sum of the items over the number of items answered. <p><u>Total Score:</u> Sum of all the items over the number of items answered on all the Scales.</p>
Interpretation and Analysis of missing data	<p>If more than 50% of the items in the scale are missing, the Scale Scores should not be computed.</p> <p>If 50% or more items are completed: Impute the mean of the completed items in a scale.</p>

PedsQL[™] Oral Health Scale

The **PedsQL™ Oral Health Scale** is composed of 5 items comprising 1 dimension.

DESCRIPTION OF THE ORAL HEALTH SCALE:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
Oral Health	5	1-5	1-5	Higher scores indicate lower problems

SCORING OF DIMENSIONS:

Item Scaling	5-point Likert scale from 0 (Never) to 4 (Almost always) 3-point scale: 0 (Not at all), 2 (Sometimes) and 4 (A lot) for the Child Report for Young Children (ages 5-7)
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed on a scale from 0 to 100.
Scoring Procedure	<p><u>Step 1: Transform Score</u></p> <p>Items are reversed scored and linearly transformed to a 0-100 scale as follows: 0=100, 1=75, 2=50, 3=25, 4=0.</p> <p><u>Step 2: Calculate Total Score</u></p> <p>Sum of all the items over the number of items answered</p>
Interpretation and Analysis of Missing Data	<p>If more than 50% of the items in the scale are missing, the Scale Score should not be computed.</p> <p>If 50% or more items are completed: Impute the mean of the completed items in a scale.</p>

PedsQL[™] Pediatric Pain Coping Inventory[™] (PPCI)

**PARENT Report for Children and Teens (ages 5-18)
CHILD Report for Children (ages 5-12), Teens (ages 13-18)**

The Parent Report of the **PedsQL™ Pediatric Pain Coping Inventory™** for Children (ages 5-12) and Teens (ages 13-18) is composed of 41 items comprising 5 dimensions.

The Child Reports of the **PedsQL™ Pediatric Pain Coping Inventory™** for:

- Children (ages 5-12)
 - And teens (ages 13-18)
- are composed of 41 items comprising 5 dimensions.

See Varni, J.W. et al. (1996). Development of the Waldron/Varni Pediatric Pain Coping Inventory. *Pain*, 67, 141-150, for scoring instructions for the *a priori* and *empirically-derived* scale scores.

DESCRIPTION OF THE PEDIATRIC PAIN COPING INVENTORY:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
Cognitive Self-Instruction	7	12, 22, 23, 24, 31, 32, 40	No	Higher scores indicate greater use of the coping strategy.
Problem-Solving	10	1, 2, 11, 15, 26, 27, 30, 33, 34, 35		
Distraction	9	6, 8, 9, 10, 14, 16, 17, 18, 19		
Seeks Social Support	9	3, 4, 7, 13, 28, 29, 37, 38, 39		
Catastrophizing / Helplessness	6	5, 20, 21, 25, 36, 41		

SCORING OF DIMENSIONS:

Item Scaling	3-point Likert scale from 0 (Never) to 2 (Often)
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed on a scale from 0 to 2.
Scoring Procedure	<u>Calculate Scale Scores</u> Sum of all the items over the number of items answered for each Scale.
Interpretation and Analysis of Missing Data	If more than 50% of the items in the scale are missing, the Scale Scores should not be computed. If 50% or more items are completed: Impute the mean of the completed items in a scale.

**PedsQL[™] Pediatric Pain Questionnaire[™]
(PPQ)**

The **PedsQL™ Pediatric Pain Questionnaire™** is composed of 3 items.

DESCRIPTION OF THE QUESTIONNAIRE:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
Present Pain	1	1	No	Higher score = higher pain intensity.
Worst Pain	1	2		

The third item refers to the localisation of pain and is not scored.

SCORING OF DIMENSIONS:

Item Scaling	10cm (100mm) VAS scales from 0 (Not hurting / No discomfort / No pain) to 10 (100) (Hurting a whole lot / Very uncomfortable / Severe pain)
Weighting of Items	No
Extension of the Scoring Scale	0-10cm or 0-100mm
Scoring Procedure	Present pain and worst pain are scored separately. The score is based on line length to the nearest 0.5 cm (5mm).
Interpretation and Analysis of Missing Data	There is no imputation process since these are single item VAS scales.

PedsQL[™] Present Functioning Visual Analogue Scale (PedsQL[™] VAS)

CHILD and PARENT reports for CHILD (ages 5-18)

The **PedsQL™ Present Functioning Visual Analogue Scale** is composed of 6 items.

DESCRIPTION OF THE SCALE:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
Present Functioning	6	1-6	No	Higher score = higher distress or intensity.

SCORING OF DIMENSIONS:

Item Scaling	100mm VAS scales from 0 to 100
Weighting of Items	No
Extension of the Scoring Scale	0-100
Scoring Procedure	<ul style="list-style-type: none"> - PedsQL VAS Total Symptom Score = average of all six items - PedsQL VAS Emotional Distress Summary Score = mean of the anxiety, sadness, anger, and worry items
Interpretation and Analysis of Missing Data	There is no imputation process since these are single item VAS scales.

PedsQL™ 3.0 Rheumatology Module

PARENT Report for Toddlers (ages 2-4)

The **Parent Report for Toddlers** (age 2-4) of the **PedsQL™ 3.0 Rheumatology Module** is composed of 14 items comprising 3 dimensions.

DESCRIPTION OF THE RHEUMATOLOGY MODULE:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
Pain and Hurt	4	1-4	1-4	Higher scores indicate lower problems.
Daily Activities	5	1-5	1-5	
Treatment	5	1-5	1-5	

SCORING OF DIMENSIONS:

Item Scaling	5-point Likert scale from 0 (Never) to 4 (Almost always)
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed to a 0 to 100 scale.
Scoring Procedure	<p><u>Step 1: Transform Score</u></p> <p>Items are reversed scored and linearly transformed to a 0-100 scale as follows: 0=100, 1=75, 2=50, 3=25, 4=0.</p> <p><u>Step 2: Calculate Scores by Dimensions</u></p> <ul style="list-style-type: none"> • If more than 50% of the items in the scale are missing, the scale scores should not be computed, • Mean score = Sum of the items over the number of items answered. <p>There is no Total Score</p>
Interpretation and Analysis of Missing Data	<p>If more than 50% of the items in the scale are missing, the Scale Scores should not be computed.</p> <p>If 50% or more items are completed: Impute the mean of the completed items in a scale.</p>

PedsQL[™] 3.0 Rheumatology Module

CHILD and PARENT reports for Young Children (ages 5-7)

The **Child and Parent Reports** of the **PedsQL™ 3.0 Rheumatology Module** for Young Children (ages 5-7) are composed of 20 items comprising 5 dimensions.

DESCRIPTION OF THE RHEUMATOLOGY MODULE:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
Pain and Hurt	4	1-4	1-4	Higher scores indicate lower problems.
Daily Activities	5	1-5	1-5	
Treatment	5	1-5	1-5	
Worry	3	1-3	1-3	
Communication	3	1-3	1-3	

SCORING OF DIMENSIONS:

Item Scaling	5-point Likert scale from 0 (Never) to 4 (Almost always) 3-point scale: 0 (Not at all), 2 (Sometimes) and 4 (A lot) for the child report for Young Children (ages 5-7)
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed on a scale from 0 to 100.
Scoring Procedure	<p><u>Step 1: Transform Score</u> Items are reversed scored and linearly transformed to a 0-100 scale as follows: 0=100, 1=75, 2=50, 3=25, 4=0.</p> <p><u>Step 2: Calculate Scores by Dimensions</u></p> <ul style="list-style-type: none"> • If more than 50% of the items in the scale are missing, the scale scores should not be computed, • Mean score = Sum of the items over the number of items answered. <p>There is no Total Score</p>
Interpretation and Analysis of Missing Data	<p>If more than 50% of the items in the scale are missing, the Scale Scores should not be computed.</p> <p>If 50% or more items are completed: Impute the mean of the completed items in a scale.</p>

PedsQL[™] 3.0 Rheumatology Module

**CHILD and PARENT reports for Children (ages 8-12) and Teens
(ages 13-18)**

The **Child and Parent Reports** of the **PedsQL™ 3.0 Rheumatology Module** for:

- Children (ages 8-12),
- And Teens (ages 13-18)

are composed of 22 items comprising 5 dimensions.

DESCRIPTION OF THE RHEUMATOLOGY MODULE:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
Pain and Hurt	4	1-4	1-4	Higher scores indicate lower problems.
Daily Activities	5	1-5	1-5	
Treatment	7	1-5	1-5	
Worry	3	1-3	1-3	
Communication	3	1-3	1-3	

SCORING OF DIMENSIONS:

Item Scaling	5-point Likert scale from 0 (Never) to 4 (Almost always)
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed on a scale from 0 to 100.
Scoring Procedure	<p><u>Step 1: Transform Score</u> Items are reversed scored and linearly transformed to a 0-100 scale as follows: 0=100, 1=75, 2=50, 3=25, 4=0.</p> <p><u>Step 2: Calculate Scores by Dimensions</u></p> <ul style="list-style-type: none"> • If more than 50% of the items in the scale are missing, the scale scores should not be computed, • Mean score = Sum of the items over the number of items answered. <p>There is no Total Score</p>
Interpretation and Analysis of Missing Data	<p>If more than 50% of the items in the scale are missing, the Scale Scores should not be computed.</p> <p>If 50% or more items are completed: Impute the mean of the completed items in a scale.</p>

PedsQL[™] 3.0 Sickle Cell Disease Module

PARENT Report for Toddlers (ages 2-4)

The Parent Report for Toddlers (age 2-4) of the PedsQL™ 3.0 Sickle Cell Disease Module is composed of 42 items comprising 9 dimensions.

DESCRIPTION OF THE SICKLE CELL DISEASE MODULE:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
Pain and Hurt	9	1-9	1-9	Higher scores indicate lower problems.
Pain Impact	10	1-10	1-10	
Pain Management	2	1-2	1-2	
Worry I	5	1-5	1-5	
Worry II	2	1-2	1-2	
Emotions	2	1-2	1-2	
Treatment	6	1-6	1-6	
Communication I	3	1-3	1-3	
Communication II	3	1-3	1-3	

SCORING OF DIMENSIONS:

Item Scaling	5-point Likert scale from 0 (Never) to 4 (Almost always)
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed to a 0 to 100 scale.
Scoring Procedure	<p><u>Step 1: Transform Score</u></p> <p>Items are reversed scored and linearly transformed to a 0-100 scale as follows: 0=100, 1=75, 2=50, 3=25, 4=0.</p> <p><u>Step 2: Calculate scores by Dimensions</u></p> <ul style="list-style-type: none"> • If more than 50% of the items in the scale are missing, the scale scores should not be computed. • Mean score = Sum of the items over the number of items answered. <p><u>Total Score:</u> Sum of all the items over the number of items answered on all the Scales.</p>
Interpretation and Analysis of Missing Data	<p>If more than 50% of the items in the scale are missing, the Scale Scores should not be computed.</p> <p>If 50% or more items are completed: Impute the mean of the completed items in a scale.</p>

PedsQL™ 3.0 Sickle Cell Disease Module

CHILD and PARENT Reports for Young Children (ages 5-7)

The **Child and Parent Reports for Young Children (ages 5-7)** of the **PedsQL™ 3.0 Sickle Cell Disease Module** are composed of 42 items comprising 9 dimensions.

DESCRIPTION OF THE SICKLE CELL DISEASE MODULE:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
Pain and Hurt	9	1-9	1-9	Higher scores indicate lower problems.
Pain Impact	10	1-10	1-10	
Pain Management	2	1-2	1-2	
Worry I	5	1-5	1-5	
Worry II*	2	1-2	1-2	
Emotions	2	1-2	1-2	
Treatment	6	1-6	1-6	
Communication I	3	1-3	1-3	
Communication II	3	1-3	1-3	

*Note: Worry II Scale is not included for child self-report for ages 5-7.

SCORING OF DIMENSIONS:

Item Scaling	5-point Likert scale from 0 (Never) to 4 (Almost always) 3-point scale: 0 (Not at all), 2 (Sometimes) and 4 (A lot) for the Child Report for Young Children (ages 5-7)
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed on a scale from 0 to 100.
Scoring Procedure	<p><u>Step 1: Transform Score</u></p> <p>Items are reversed scored and linearly transformed to a 0-100 scale as follows: 0=100, 1=75, 2=50, 3=25, 4=0</p> <p><u>Step 2: Calculate Scores by Dimensions</u></p> <ul style="list-style-type: none"> • If more than 50% of the items in the scale are missing, the scale scores should not be computed. • Mean score = Sum of the items over the number of items answered. <p><u>Total Score:</u> Sum of all the items over the number of items answered on all the Scales.</p>
Interpretation and Analysis of Missing Data	<p>If more than 50% of the items in the scale are missing, the Scale Scores should not be computed.</p> <p>If 50% or more items are completed: Impute the mean of the completed items in a scale.</p>

PedsQL™ 3.0 Sickle Cell Disease Module

**CHILD and PARENT Reports for Children (ages 8-12) and
Teens (ages 13-18)**

The **Child and Parent Reports for Children (ages 8-12) and Teens (ages 13-18)** of the **PedsQL™ 3.0 Sickle Cell Disease Module** are composed of 43 items comprising 9 dimensions.

DESCRIPTION OF THE SICKLE CELL DISEASE MODULE:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
Pain and Hurt	9	1-9	1-9	Higher scores indicate lower problems.
Pain Impact	10	1-10	1-10	
Pain Management	2	1-2	1-2	
Worry I	5	1-5	1-5	
Worry II	2	1-2	1-2	
Emotions	2	1-2	1-2	
Treatment	7	1-7	1-7	
Communication I	3	1-3	1-3	
Communication II	3	1-3	1-3	

SCORING OF DIMENSIONS:

Item Scaling	5-point Likert scale from 0 (Never) to 4 (Almost always)
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed on a scale from 0 to 100.
Scoring Procedure	<p><u>Step 1: Transform Score</u></p> <p>Items are reversed scored and linearly transformed to a 0-100 scale as follows: 0=100, 1=75, 2=50, 3=25, 4=0</p> <p><u>Step 2: Calculate Scores by Dimensions</u></p> <ul style="list-style-type: none"> • If more than 50% of the items in the scale are missing, the scale scores should not be computed. • Mean score = Sum of the items over the number of items answered. <p><u>Total Score:</u> Sum of all the items over the number of items answered on all the Scales.</p>
Interpretation and Analysis of Missing Data	<p>If more than 50% of the items in the scale are missing, the Scale Scores should not be computed.</p> <p>If 50% or more items are completed: Impute the mean of the completed items in a scale.</p>

PedsQL[™] 3.0 Transplant Module

The **PedsQL™ 3.0 Transplant Module** is composed of 46 items comprising 8 dimensions.

DESCRIPTION OF THE TRANSPLANT MODULE:

Dimensions	Number of Items	Cluster of Items	Reversed Scoring	Direction of Dimensions
About My Medicines I	9	1-9	1-9	Higher scores indicate lower problems.
About My Medicines II	8	1-8	1-8	
My Transplant and Others	8	1-8	1-8	
Pain and Hurt	3	1-3	1-3	
Worry	7	1-7	1-7	
Treatment Anxiety	4	1-4	1-4	
How I Look	3	1-3	1-3	
Communication	4	1-4	1-4	

SCORING OF DIMENSIONS:

Item Scaling	5-point scale from 0 (Never) to 4 (Almost always) 3-point scale: 0 (Not at all), 2 (Sometimes) and 4 (A lot) for the Child Report for Young Children (ages 5-7)
Weighting of Items	No
Extension of the Scoring Scale	Scores are transformed on a scale from 0 to 100.
Scoring Procedure	<p><u>Step 1: Transform Score</u></p> <p>Items are reversed scored and linearly transformed to a 0-100 scale as follows: 0=100, 1=75, 2=50, 3=25, 4=0.</p> <p><u>Step 2: Calculate scores by Dimension</u></p> <ul style="list-style-type: none"> • If more than 50% of the items in the scale are missing, the scale scores should not be computed, • Mean score = Sum of the items over the number of items answered. <p><u>Total Score:</u> Sum of all the items over the number of items answered on all the Scales.</p>
Interpretation and Analysis of Missing Data	<p>If more than 50% of the items in the scale are missing, the Scale Scores should not be computed.</p> <p>If 50% or more items are completed: Impute the mean of the completed items in a scale.</p>